

Synergy in Action: Microbial Consortia for Environmental Clean-Up

Abisha R. J., Shri Vasanthan K. K., Arasi. R., D. Manimekalai and V. Rani.

Department of Aquatic Environment Management, Tamil Nadu Dr. J. Jayalalithaa Fisheries University,
Fisheries College and Research Institute, Thoothukudi, Tamil Nadu, India

SUMMARY

Microbial consortia, composed of diverse groups of microorganisms working synergistically, offer a promising approach to environmental protection. These consortia can be harnessed for bioremediation, wastewater treatment, soil restoration, and pollutant degradation. By leveraging the complementary metabolic capabilities of multiple microbial species, consortia enhance the breakdown of hazardous substances, including heavy metals, organic pollutants, and nutrients, in various environmental settings. Their use in enhancing soil fertility, mitigating water pollution, and promoting carbon sequestration makes them a sustainable and efficient solution for addressing environmental challenges. The adaptability and resilience of microbial consortia under varying conditions further contribute to their potential as a green technology for restoring and protecting ecosystems.

INTRODUCTION

In the natural environment, harmful pollutants can be removed through biological, chemical, and photochemical degradation. Physical and chemical methods of removing pollutants include coagulation, precipitation, centrifugation, adsorption and desorption, hydrolysis, and photodegradation and these processes are costly and have limited effects. Microbial consortia consist of multiple microbial species, including bacteria, fungi, archaea, and algae, that coexist and interact within a specific environment. These interactions can be synergistic, where the presence of multiple species enhances overall functionality, or competitive, influencing community dynamics and stability. The microbial consortium has become an important technology because it degrades pollutants more effectively than a single strain. Bioremediation is usually carried out by the microbial consortium rather than by individual species in the natural environment and different strains or species play different functional roles.

Concept of microbial consortia:

Microbial consortia functions based on the interspecies cooperation, where the metabolic by-products of one species can serve as substrates for others, facilitating more complete degradation or transformation of pollutants. This cooperation often leads to enhanced resilience and efficiency compared to single-species systems. These consortia can be naturally occurring or engineered to perform specific functions. These communities can degrade pollutants, recycle nutrients, or transform toxic substances into less harmful compounds.

Synergy: Different microbes often have complementary metabolic capabilities. For example, one microbe might break down a complex pollutant into simpler compounds, which another microbe can further degrade.

Stability and Resilience: Diverse microbial communities are often more stable and resilient to environmental changes than single-species systems, which is crucial for consistent performance in dynamic environments. The microbial consortia can be adapted more rapidly to environmental changes than monocultures. By providing metabolic versatility, the consortia improve system stability, crucial for large-scale environmental interventions like bioremediation and wastewater treatment.


Mechanisms of bioremediation of contaminated sites by microbial consortium:

- The bacterial consortium members can degrade the essential intermediate compounds produced by other members in degrading pollutants and reducing the accumulation of intermediate products, thereby increasing the synergistic metabolic degradation.
- Some bacterial consortium strains produce many high-efficiency biosurfactants, thereby increasing the solubility and content of pollutants, improving their bioavailability and biodegradability.
- The microbial consortium can self-regulate and adapt during degradation. Microbial consortia show better performance than individual cultures in degrading contaminants.

- The microbial consortium can promote the growth of strains by using metabolites after pollutant degradation.
- The crude enzyme produced in the microbial consortium's intracellular space can be used as a degradation factor in degradation, showing high degradation activity.
- There is a biochemical synergistic effect between bacterial consortium strains, which enhances bacterial activity and the degradation of pollutants.

Example: Efficient removal of tetracycline (TC) by bacterial consortia:

Tetracycline (TC), an environmental pollutant, can stay in the soil for many years and can destroy the ecosystem. The TC-degrading bacterial consortium composed of *Raoultella* sp. XY-1 and *Pandoraea* sp. XY-2 strains that was isolated and constructed from TC-contaminated soil. Compared to a single strain, this TC-degrading bacterial consortium grew better and was able to degrade TC more efficiently. This is due to the presence of a biochemical synergistic effect between the bacteria, which enhanced the bacteria's activity and the TC degradation.



Applications of microbial consortia:

Bioremediation:

Microbial consortia are widely employed to degrade or detoxify environmental pollutants like hydrocarbons, heavy metals, pesticides, and other toxic compounds. The diverse metabolic pathways of different microbes in the consortium allow for the breakdown of complex substances that single species may not effectively handle. Microbial consortia can be used to degrade oil spills, heavy metals, pesticides, or other hazardous substances. For example, *Pseudomonas*, *Bacillus*, and *Acinetobacter* species are often used to degrade oil spills. *Lactobacilli*, *Actinobacteria*, *Pseudomonas*, *Clostridium*, *Salmonella*, and *Escherichia coli* have been found to have the inherent ability to degrade pollutants.

Hydrocarbon degradation:

Hydrocarbons, such as those found in petroleum products, are major environmental pollutants, particularly in oil spills, contaminated soils, and groundwater. The consortium of *Pseudomonas putida*, *Bacillus subtilis*, and *Rhodococcus* sp. effectively degraded polycyclic aromatic hydrocarbons (PAHs) in contaminated soil, achieving significant reduction in pollutant levels. In the presence of oxygen, the consortia consisting of species like *Pseudomonas*, *Alcanivorax*, and *Rhodococcus* are effective at degrading hydrocarbons such as alkanes, Polycyclic Aromatic Hydrocarbons (PAHs), and aromatic compounds. These microbes produce enzymes such as oxygenases and hydroxylases that initiate the oxidation of hydrocarbons into alcohols, aldehydes, and carboxylic acids, which are further broken down into CO₂ and water. In oxygen-deprived environments like oil reservoirs and deep-sea sediments, the anaerobic microbial consortia which includes the bacteria from the genera *Desulfobacter*, *Geobacter*, and *Syntrophus* that utilize alternate electron acceptors like nitrate, sulfate, or iron to degrade hydrocarbons.

Heavy metal removal:

Heavy metals such as lead (Pb), cadmium (Cd), chromium (Cr), and arsenic (As) are persistent pollutants that pose serious risks to ecosystems and human health. Microbial consortia can facilitate the removal of these metals through various mechanisms, including biosorption, bioprecipitation, and bioaccumulation. Microbial cell walls, primarily composed of polysaccharides, proteins, and lipids, contain functional groups (e.g., carboxyl, hydroxyl, and phosphate) that can bind heavy metals from contaminated environments. The consortia, such as those containing *Bacillus*, *Pseudomonas*, and *Aspergillus*, efficiently adsorb metals on their surfaces. Certain microbial consortia can transform soluble metal ions into insoluble forms through enzymatic processes, facilitating their precipitation out of solution. For example, sulfate-reducing bacteria like *Desulfovibrio* reduce sulfate to hydrogen sulfide, which reacts with metals like zinc and cadmium to form insoluble metal sulfides. For example, *Geobacter* species can reduce hexavalent chromium (Cr^{6+}) to its less toxic trivalent form (Cr^{3+}), and *Shewanella* species can reduce soluble uranium (U^{6+}) to insoluble U^{4+} , effectively immobilizing it.

Wastewater treatment:

In wastewater treatment, microbial consortia helps in the removal of organic matter, nitrogen, and phosphorus through processes like nitrification, denitrification, and aerobic/anaerobic digestion. Microbial consortia are integral to the functioning of wastewater treatment plants (WWTPs), facilitating the removal of organic matter, nitrogen, and phosphorus. Mixed cultures, such as ammonia-oxidizing bacteria and methanogens, work together to stabilize waste products, reducing pollutants and producing methane that can be used for energy. For example, sulfate-reducing bacteria (*Desulfovibrio* sp.) combined with *Bacillus* species has the enhanced potential for the removal of cadmium and lead from wastewater through sulfide precipitation.

Scope of microbial consortia for environmental protection:**Ecosystem restoration:**

Microbial consortia play a crucial role in restoring degraded ecosystems by re-establishing essential microbial processes and facilitating the recovery of native flora and fauna. Microbial consortia have been applied in restoring polluted marine environments, such as those affected by oil spills, and in soil remediation. Microbial communities, through natural processes, can enhance soil fertility and break down hydrocarbons in marine ecosystems, facilitating ecological recovery.

Microbial Fuel Cells (MFCs):

Consortia of electroactive bacteria can be harnessed in microbial fuel cells (MFCs) for bioremediation and electricity generation. In these systems, microbes metabolize organic waste and release electrons, producing electricity while simultaneously cleaning the waste.

Carbon sequestration and climate change mitigation:

Microbial consortia in soil and oceans play a vital role in carbon cycling and sequestration. It contributes to climate change mitigation by sequestering carbon dioxide and reducing greenhouse gas (GHG) emissions. For example, the consortia of methanotrophs and methanogens can effectively sequester carbon dioxide and mitigated methane emissions from agricultural soils.

Agriculture:

In agriculture, microbial consortia improve soil health, enhance nutrient availability, and promote plant growth through natural processes. For example, the consortium comprising nitrogen-fixing bacteria, phosphate-solubilizing bacteria, and plant growth-promoting rhizobacteria (PGPR) is found to be significantly enhancing crop yields and soil fertility.

Feasibility of using microbial consortia for environmental protection:

Selection and optimization: Identifying the right combination of microorganisms and optimizing their interactions is critical for achieving the desired environmental outcomes. Advances in metagenomics, synthetic biology, and systems biology are aiding in the design and optimization of microbial consortia.

Scalability: Implementing microbial consortia at a large scale requires careful consideration of factors such as nutrient availability, environmental conditions, and the potential impact on native microbial communities.

Regulation and safety: The release of engineered microbial consortia into the environment raises regulatory and safety concerns. It is essential to assess the potential risks and ensure that the introduced microbes do not negatively impact native species or ecosystems.

Cost-effectiveness: While microbial consortia offer a potentially sustainable solution, the costs associated with developing, deploying, and monitoring these consortia must be considered.

Monitoring and control: Continuous monitoring of microbial consortia and their environmental impacts is crucial for ensuring they perform as expected. Strategies for controlling the consortia, such as using kill switches in engineered microbes, may be necessary to prevent unintended consequences.

CONCLUSION

Microbial consortia offer significant potential for tackling a range of environmental issues, including pollution, wastewater treatment, bioremediation, and climate change mitigation. Their ability to interact and cooperate makes them powerful tools for maintaining ecological balance and protecting environmental health. While microbial consortia offer great potential, challenges include maintaining stability and activity in varying environmental conditions and avoiding competition between microbial species. Ongoing research focuses on genetically engineering or selecting optimal microbial consortia for enhanced performance in specific contaminated environments. In summary, microbial consortia play a crucial role in the bioremediation of hydrocarbons and heavy metals, leveraging the metabolic diversity and cooperative interactions between different microbial species to achieve efficient environmental cleanup.

REFERENCES

- Brahmaprakash, G.P., Sahu, P.K., Lavanya, G., Nair, S.S., Gangaraddi, V.K. and Gupta, A., 2017. Microbial functions of the rhizosphere. *Plant-Microbe Interactions in Agro-Ecological Perspectives: Volume 1: Fundamental Mechanisms, Methods and Functions*, pp.177-210.
- Jain, A., Chatterjee, A. and Das, S., 2020. Synergistic consortium of beneficial microorganisms in rice rhizosphere promotes host defense to blight-causing *Xanthomonas oryzae* pv. *oryzae*. *Planta*, 252(6), p.106.
- Negi, R., Sharma, B., Jan, T., Kaur, T., Chowdhury, S., Kapoor, M., Singh, S., Kumar, A., Rai, A.K., Rustagi, S. and Shreaz, S., 2024. Microbial Consortia: Promising Tool as Plant Bioinoculants for Agricultural Sustainability. *Current Microbiology*, 81(8), p.222.
- Niu, B., Wang, W., Yuan, Z., Sederoff, R.R., Sederoff, H., Chiang, V.L. and Borriss, R., 2020. Microbial interactions within multiple-strain biological control agents impact soil-borne plant disease. *Frontiers in Microbiology*, 11, p.585404.
- Uddin, T.M., Chakraborty, A.J., Khusro, A., Zidan, B.R.M., Mitra, S., Emran, T.B., Dhama, K., Ripon, M.K.H., Gajdács, M., Sahibzada, M.U.K. and Hossain, M.J., 2021. Antibiotic resistance in microbes: History, mechanisms, therapeutic strategies and future prospects. *Journal of infection and public health*, 14(12), pp.1750-1766.