

Phragmosis: The Ingenious Defense Mechanism of Arthropods

Malireddi Prasanna¹, Mounika Jarpla², R. D. Patel³ and Bhanavathu Sai¹

¹M.Sc. Scholar, Department of Entomology, Navsari Agricultural University, Navsari Gujarat

²Ph.D. Scholar, Department of Entomology, Navsari Agricultural University, Navsari, Gujarat

³Assistant Research Scientist, Main Cotton research station, Surat, Navsari Agricultural University, Gujarat

SUMMARY

Phragmosis stands as a captivating defensive behavior observed across diverse arthropod species, ranging from insects to arachnids, demonstrating the ingenious strategies these creatures employ to protect themselves from predators and environmental adversities. This adaptive response entails the deliberate blocking or sealing of entrances to their shelters or hiding spots, often leveraging specialized anatomical features or external materials for the task. The phenomenon's species-specific adaptations underscore the rich variety of antipredator tactics existing in nature, where each species may manifest unique morphological attributes or behavioral patterns tailored to its ecological niche and evolutionary trajectory. For instance, certain insects might possess specialized appendages or modifications to their exoskeletons facilitating effective barricading, while select arachnids may rely on silk production to fortify their refuges. Phragmosis transcends mere physical obstruction, encompassing a spectrum of behavioral nuances shaped by evolutionary pressures and environmental cues. Arthropods may exhibit sophisticated movement patterns and coordination, demonstrating acute awareness of predator presence and the exigency for rapid defensive responses. Understanding the adaptive advantages conferred by phragmosis yields insights into predator-prey dynamics and community ecology, whereby fortified shelters can reduce predation risks and bolster survival chances, thereby shaping population dynamics and species interactions within ecosystems.

INTRODUCTION

W.M. Wheeler (1927) introduced the term "phragmosis" to characterize a secretive defensive strategy utilized by insects, wherein they utilize uniquely adapted body structures to obstruct entrances to their nests (Brandão *et al.*, 2001). "Phragmosis," derived from the Greek word (pronounced "fráɡma"), meaning "barrier" or "shield," represents an evolutionary tactic that combines specialized morphology (the shield) with corresponding behavior to protect a narrow opening (chokepoint) by an individual or a small group (Bishoff *et al.*, 2023). Phragmosis, initially associated with insects, has been expanded to include other taxa. When an animal resides in a protective nest, the entrance becomes a vulnerable point susceptible to potential attacks by predators or parasites. One defense mechanism involves the evolution of both a morphologically resilient surface, typically a hard, flat, and thick body part referred to as a "shield," that perfectly fits the entryway, along with a corresponding behavioral response. This behavior entails utilizing the shield to block the entrance when facing threats. For optimal effectiveness, the shield should lack any structures that predators can grasp onto, while potentially featuring sharp spines, dense hair, or other deterrent features to repel intruders. In insects, it's common for either the head (occasionally with thoracic nota) or the tip of the abdomen to develop a particularly hardened region. This area typically adopts a truncated shape with edges that precisely match the entrance of their nest. In response to threats, insects utilize this hardened region to effectively block the nest entrance (Bishoff *et al.*, 2023).

Phragmosis in Aphids

The aphid species *Astegopteryx sp.* constructs a gall resembling a cluster of bananas on *Styrax benzoides* trees in northern Thailand. It undergoes its entire life cycle on the same tree without migrating to other host plants. Notably, the soldiers of this aphid exhibit sclerotic, protruding heads adorned with numerous spine-like setae. These soldiers collaborate to seal the ostiole of the sub gall using their heads which helps in defense mechanism (Kurosu *et al.*, 2006).

Phragmosis in Ants

The defensive strategy of using a soldier's head to block a nest entrance is a recognized behavior observed in certain ant species (Hasegawa, 1993). Soldiers and queens of Neotropical myrmicine *Cephalotes*

species within the pallens group possess shield-like heads, seemingly secreting a fibrous substance from numerous pores reminiscent of glandular openings. This substance coats the head disk, gradually accumulating into a dense mass. Although superficially resembling fungal mycelia, it lacks the branching pattern and internal structure characteristic of fungi (De Andrade & Urbani, 1999). In contrast to *Cephalotes*, other ant genera develop phragmotic structures through the accumulation of environmental debris held by specialized hairs (Wheeler and Hölldobler 1985). In the genus *Pheidole* the queen uses “reverse phragmosis” where a morphologically flattened posterior “plug-like modification of the gaster” defends the brood chamber (Brown, 1967).

Phragmosis in Arachnids

Certain arachnids exhibit phragmotic behavior, as seen in trapdoor spiders. These spiders construct tunnels within the ground, which they line with silk and cover tightly with a door made of silk and debris (Bishoff *et al.*, 2023). Some trapdoor spiders, in genera such as *Cyclocosmia* and *Ummidia* have a phragmotic abdomen (Xu *et al.*, 2017). If the trapdoor being breached, these arachnids rely on their truncated, robust abdomens to shield themselves from predators like birds, scorpions, and snakes within their tunnels (Bond and Coyle, 1995).

Phragmosis in Termites

Soldier castes among termites are the most highly specialized of all social insects, (Deligne *et al.*, 1981). Across most termite species, soldiers are dedicated solely to defense and lack the ability to reproduce, perform work tasks, or gather food independently. Consequently, the soldier caste assumes a pivotal role in safeguarding the colony, albeit at the expense of reproduction and self-sustenance (Matsuura, 2002). Soldiers guarded the nest openings with their heads and used their mandibles.

CONCLUSION

Phragmosis represents a remarkable defensive strategy observed across diverse arthropod and amphibian species, showcasing the ingenious adaptations organisms have evolved to protect themselves from predators and environmental challenges. This behavior involves the deliberate blocking or sealing of entrances to shelters, often leveraging specialized anatomical features or external materials. The diversity of phragmotic adaptations underscores the rich variety of antipredator tactics in nature, with each species manifesting unique morphological attributes or behavioral patterns tailored to its ecological niche. From insects to arachnids, ants to termites, and even frogs, phragmosis manifests in various forms, each tailored to the specific needs and habitats of the organism. Through detailed studies and observations, researchers have gained insights into the evolutionary origins, ecological significance, and adaptive advantages conferred by phragmosis. Understanding these defensive mechanisms not only enriches our understanding of animal behavior but also sheds light on the complex dynamics of predator-prey interactions and community ecology. Moreover, the study of phragmosis offers valuable insights into the broader principles of adaptation and resilience, with potential implications for conservation and management efforts aimed at preserving biodiversity and ecosystem integrity.

REFERENCES

- Bishoff, M. J., Peng, L., Zang, H. M., & Morse, J. C. (2023). Defensive phragmosis and cathaptosis in Trichoptera larvae. *Contributions to Entomology*, 73(2), 209-218.
- Bond, J. E., & Coyle, F. A. (1995). Observations on the natural history of an *Ummidia* trapdoor spider from Costa Rica (Araneae, Ctenizidae). *Journal of Arachnology*, 157-164.
- Brandão, C. R. F., Diniz, J. L., Silva, P. R., Albuquerque, N. L., & Silvestre, R. (2001). The first case of intranidal phragmosis in ants. The ergatoid queen of *Blepharidatta conops* (Formicidae, Myrmicinae) blocks the entrance of the brood chamber. *Insectes Sociaux*, 48, 251-258.
- Brown, W. L. (1967). A New *Pheidole* with Reversed Phragmosis (Hymenoptera: Formicidae). *Psyche: A Journal of Entomology*, 74(4), 331-339.
- De Andrade, M. L., & Urbani, C. B. (1999). Diversity and adaptation in the ant genus *Cephalotes*, past and present. *Staatliches Museum für Naturkunde*.
- Deligne, J. E. A. N., Quennedey, A. N. D. R. E., & Blum, M. S. (1981). The enemies and defense mechanisms of termites. *Social insects*, 2, 1-76.

- Hasegawa, E. (1993). Nest defense and early production of the major workers in the dimorphic ant *Colobopsis nipponicus* (Wheeler) (Hymenoptera: Formicidae). *Behavioral ecology and sociobiology*, 33, 73-77.
- Kurosu, U., Narukawa, J., Buranapanichpan, S., & Aoki, S. (2006). Head-plug defense in a gall aphid. *Insectes sociaux*, 53, 86-91.
- Matsuura, K. (2002). Colony-level stabilization of soldier head width for head-plug defense in the termite *Reticulitermes speratus* (Isoptera: Rhinotermitidae). *Behavioral Ecology and Sociobiology*, 51, 172-179.
- Wheeler, D. E., & Hölldobler, B. (1985). Cryptic phragmosis: the structural modifications. *Psyche: A Journal of Entomology*, 92, 337-353.
- Xu, X., Xu, C., Li, F., Pham, D. S., & Li, D. Q. (2017). Trapdoor spiders of the genus *Cyclocosmia* Ausserer, 1871 from China and Vietnam (Araneae, Ctenizidae). *ZooKeys* 643: 75-85.